

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
28 October 2004 (28.10.2004)

PCT

(10) International Publication Number
WO 2004/091314 A2

(51) International Patent Classification⁷: **A23L 1/20**,
1/10, 1/36, A23C 11/10, A23L 1/03

(21) International Application Number:
PCT/US2004/009366

(22) International Filing Date: 26 March 2004 (26.03.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/458,943 1 April 2003 (01.04.2003) US
60/519,795 14 November 2003 (14.11.2003) US

(71) Applicant (for all designated States except US):
ARCHER-DANIELS-MIDLAND COMPANY
[US/US]; 4666 East Faries Parkway, Decatur, IL 62526
(US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **GOTTEMOLLER,**
Thomas [US/US]; 1565 Alexander Drive, Mt. Zion, IL
62549 (US).

(74) Agents: **HIRSHMAN, Jesse, A. et al.**; Kirkpatrick &
Lockhart LLP, Henry W. Oliver Building, 535 Smithfield
Street, Pittsburgh, PA 15222-2312 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,
GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **SOYA FIBER PARTICULATES AND METHODS OF PREPARATION**

(57) Abstract: The present invention belongs to the field of processing techniques used in the manufacture of soyfoods. The inven-
tion relates to methods for preparing a soya fiber particulate from dehulled soybeans of a superior smooth texture, nutritional quality
and flavor. The invention can be used to obtain a soya fiber particulate having a particle size in a range of about 0.01 microns to
about 100 microns, wherein at least about 50% to about 100% of the soya fiber particulate has a size in a range of about 0.01 microns
to about 35 microns. The invention furthermore is concerned with the use in the manufacture of food products which include but are
not limited to beverages, dry mixes, yogurts, sour cream, and cream cheese.

WO 2004/091314 A2

SOYA FIBER PARTICULATES AND METHODS OF PREPARATION**INVENTOR**

Thomas Gottemoller

BACKGROUND OF THE INVENTION5 **1. Field of the Invention**

The present invention is related to the field of processing techniques used in the manufacture of soyfoods. The invention provides methods for preparing a soya fiber particulate from dehulled soybeans. The invention can be used to obtain a soya fiber particulate having a particle size in a range of about 0.01 microns to about
10 100 microns, wherein at least about 50% to about 100% of the soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns. The soya fiber particulate may be used alone or as an ingredient in food products.

2. Related Art

The soybean is the one of the few plant protein sources that provides
15 all eight essential amino acids. Soy protein is of the highest quality, equal to that of meat and dairy products, but without the cholesterol and high levels of saturated fat. In addition, the soybean has a higher amount of protein than other beans. Research has revealed many possible health benefits that may be achieved from the consumption of food products containing soy protein. The bioactive molecules
20 occurring naturally in soy protein have antioxidant activities, which enhance the immune system, the body's defense against a wide range of diseases. For example, soy protein may play a role in the reduction of the risk of developing cardiovascular disease, including preventing heart attack, stroke, and high blood pressure. Soy protein, which is contained within soya fiber particulates, may also play a role in the

reduction of the risk of developing kidney disease, various cancers and diabetes. Soy protein may also help protect against the development of osteoporosis and promote weight management. Because of these and other potential health effects, the commercial demand for soya fiber particulates is expected to rise rapidly.

5 Soymilk, an aqueous extraction of soybeans, is used as a base in a wide variety of products. There are already a few reported processing techniques used in the manufacture of soymilk. A traditional method for soymilk production involves cleaning and then soaking whole, undehulled soybeans in cold water, rinsing the soaked soybeans, grinding the rinsed soybeans with additional fresh water into a
10 coarse slurry, heating and then centrifuging the slurry, retaining the soluble soymilk fraction and disposing of the okara residue, also described as soymilk pulp or cell wall debris.

One disadvantage of traditional methods for soymilk production is the presence of an undesirable beany flavor, odor, and color. In order to improve the
15 flavor of soymilk, alternate methods include heating and/or blanching for the inactivation of the enzymes that are responsible for the undesirable beany flavor and subsequent extraction of soluble constituents. For example, U.S. Patent No. 4,039,696 describes a process for preparing a milk-like product from whole soybeans by hot aqueous alkaline blanching and centrifugation. U.S. Patent No. 4,971,825
20 describes a process for producing soybean milk which involves crushing, rather than grinding, soybeans in aqueous hot medium followed by soy protein extraction. U.S. Patent No. 4,992,294 discloses a method of producing soybean milk by boiling soybeans and straining the boiled solution. U.S. Patent No. 5,945,151 describes a process for producing a soymilk which involves heating a soybean paste. U.S. Patent

No. 4,186,218 describes a process of improving denatured soy protein by superheating and/or alcohol extraction.

Other methods for eliminating undesirable odors from soymilk include fermentation. For example, U.S. Patent No. 3,971,843 describes a process for
5 eliminating a bean odor from a soymilk by fermentation and distillation under reduced pressure. U.S. Patent No. 4,664,919 describes a process for producing a liquid-like lactic acid soymilk drinks by inoculating filtered pulverized soybeans with *Streptococcus sojilactis* bacterium.

Soymilk produced according to traditional methods has a grainy
10 texture. Occasionally, this grainy texture is desirable. For example, U.S. Patent No. 6,413,557 describes a process for preparing a textured soya beverage which includes blending soaked, rinsed, and cooked soybeans with loose soybean hulls. However, a soymilk composition with a smooth texture and pleasant mouthfeel is usually preferable. In order to obtain a soymilk with a smooth texture, alternate methods
15 include the addition of enzymes to soybeans to degrade proteinaceous, carbohydrate, and cellulosic constituents, preventing sedimentation. For example, U.S. Patent No. 3,941,890 describes a process for preparing soymilk from dehulled soybeans by microwaving and adding suitable enzymatic material. U.S. Patent No. 4,378,376 describes a process for producing a simulated milk protein product by forming a
20 neutralized aqueous slurry of a soya protein material and dairy whey and then reacting the slurry with a proteolytic enzyme.

Other methods for obtaining a soymilk with a smooth texture include multiple or extensive grinding steps. For example, U.S. Patent No. 5,807,602 describes a process for making a finely divided soya product based on continuous
25 milling, superheating and flash evaporation. This method obtains soya particles

wherein 90% have a diameter of less than 85 microns and 50% have a diameter of less than 40 microns. U.S. Patent No. 4,194,018 describes a process of preparing a soya composition from soybeans yielding particulate in which the largest dimension does not exceed 40 to 300 microns.

5 Finally, another disadvantage of traditional methods for soymilk production is that the full nutritional quality of the soybean is not maximized. In order to improve the nutritional quality of soymilk, alternate methods include the use of acid or acid salt to process soybeans; specifically to chelate various soybean components. For example, U.S. Patent No. 6,322,846 describes a process for
10 preparing soymilk and beverage compositions from whole soybeans which involves the addition of at least one of an acid and/or an acid salt. U.S. Patent No. 4,389,425 describes a process for preparing a bovine milk substitute which includes grinding soybeans in the presence of an acidified water solution followed by homogenization, desludging, and extraction.

15 Other methods for improving the nutritional quality of soymilk involve extraction and/or filtration. For example, U.S. Patent No. 3,995,071 describes a process based on the aqueous extraction of defatted soy flakes, basefication, and the removal of insolubles. U.S. Patent No. 4,091,120 describes a process for preparing soya protein beverages by ultrafiltration purification and precipitation.

20 All of these reported processes are unable to produce a soya fiber particulate that has, simultaneously, a very smooth texture, a high nutritional value and lacks a beany flavor. For example, while the process disclosed in U.S. Patent No. 4,194,018 produces a soya composition of a somewhat improved texture, the largest dimension of the product does not exceed 50 to 300 microns, meaning that the
25 composition still has a grainy mouthfeel and lacks the stability for the prevention of

sedimentation of the soy fiber particulate during storage. Therefore, there is a great need for improved, economical processes that provide very smooth soya fiber particulates.

SUMMARY OF THE INVENTION

5 The invention is based on the surprising fact that soybeans can be processed such that a soya fiber particulate having a particulate size in a range of about 0.01 microns to about 100 microns is obtained. The soya fiber particulate of the invention has a very smooth texture, because about 50% to about 100% of the soya fiber particulate has a particle size in a range of about 0.01 microns to about 35
10 microns. The invention utilizes, rather than disposes of, the okara, which contains high levels of fiber and also comprises iron, calcium, protein, carbohydrate, thiamin, riboflavin, and niacin. Therefore, in addition to its smooth texture, the soya fiber particulate of the present invention is of high nutritional quality. This utilization of the okara by the processes of the present invention is highly economical, due to the
15 high disposal costs of okara incurred by other processes. Finally, the soya fiber particulate of the present invention lacks a beany flavor and has a high stability. The technology allows for the use of this soya fiber particulate alone or as an ingredient in food products.

 It is to be understood that both the foregoing general description and
20 the following detailed description are exemplary and explanatory only and are intended to provide further explanation of the invention as claimed.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

 Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to

which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are hereinafter described.

1. Definitions

5 Each of the foregoing definitions is illustrative, and is not intended to be limiting.

Beverage: As used herein, the term "beverage" is taken to include, for example, energy drink, sports drink, coffee, beverage powder, liquid soymilk, powdered soymilk, soymilk and milk replacer.

10 Cotyledon: As used herein, the term "cotyledon" means the usable meat of the soybean. The complex carbohydrate component of the soybean is contained in the cell walls of the cotyledon.

Dehulled: As used herein, the term "dehulled" means the hulls of the soybeans, which are the outer layer of cells which coat the cotyledons, have been
15 removed. The hulls, which are highly cellulosic and make up about 8% of the bean's total weight, may be removed from soybeans by mechanical means, for example, by cracking or crushing. The hulls, which have been removed from dry cracked soybeans, may be separated from cotyledons by blowing. Alternatively, hulls may be removed from cracked soybeans by exposure to heat, for example, to boiling water.
20 The hulls, which have been removed from boiled soybeans, may be washed away from the cotyledons by a stream of water and/or action of a stationary or a moving object, for example, a brush.

Dehydrated: As used herein, the term "dehydrated" means to remove water from or to make anhydrous.

Dessert: As used herein, the term "dessert" is taken to include, for example, cakes, sponge cakes, wafers, confections, pudding, cookies, sweet pastries, fillings, muffins, biscuits and frozen desserts.

Dry mix: As used herein, the term "dry mix" is taken to include, for example, flakes, breakfast cereal, soup mix, falafel mix, pancake mix and cake mix.

Flour product: As used herein, the term "flour product" is taken to include, for example, soy flour, breads, for example, white bread and multigrain bread, buns, rolls, waffles, pancakes and pasta products.

Grind: As used herein, the term "grind" is taken to include, for example, chop, crumble, crush, disintegrate, granulate, grate, mill, pestle, pound, powder and pulverize.

Infant food: As used herein, the term "infant food" is taken to include, for example, infant formula.

Meat food product: As used herein, the term "meat food product" is taken to include, for example, bologna, frankfurters, burgers, sausage, luncheon, loaves, bacon, and meatballs. A meat food product may be combined with a simulated meat food product.

Milk product: As used herein, the term "milk product" is taken to include, for example, a soymilk product, sour cream, ice cream, whipped cream, yogurt, cheese, coffee whitener, coffee creamer and cream cheese. A milk product may be combined with a simulated milk product.

One, a or an: When the terms "one," "a," or "an" are used in this disclosure, they mean "at least one" or "one or more," unless otherwise indicated.

Simulated meat food product: As used herein, the term "simulated meat food product" is taken to include, for example, a meat alternative, meat analog,

soy burger, soy bologna, soy frankfurter, soy sausage, soy luncheon loaves, soy bacon and soy meatball. A simulated meat food product may be combined with a meat food product.

Simulated milk product: As used herein, the term "simulated milk product" is taken to include, for example, a milk alternative, milk analog, soymilk product, soy sour cream, soy ice cream, soy whipped cream, soy yogurt, soy cheese, soy coffee whitener, soy coffee creamer and soy cream cheese. A simulated milk product may be combined with a milk product.

Snack: As used herein, the term "snack" is taken to include, for example, energy bar, nutrition bar, protein bar, crackers, doughnuts and chips. The term "snack" is also taken to include extruded snacks, for example, cheese balls, cheese curls, and rice pieces.

Soya fiber particulate: As used herein, the term "soya fiber particulate" is taken to mean a suspension of ground soya. Suitable soya include but is not limited to whole soybeans, full fat soy flour, full fat soy flakes, partially defatted soy flour, partially defatted soy flakes, defatted soy flour, defatted soy flakes, refatted soy flour, refatted soy flakes, soy protein concentrate and mixtures thereof. The ground soya may be suspended in water or a water-based product. The soya fiber particulate may include, for example, finely ground okara and cell wall debris. The soya fiber particulate may be concentrated into a powder form by evaporation or by dehydration. Alternatively, the soya fiber particulate may remain in a liquid form.

Spray drying: As used herein, the term "spray drying" means transforming a fluid medium into a dry-powdered or particulate form. This is achieved by atomizing the fluid into a drying chamber, where the liquid droplets are passed through a hot-air stream. The objective is to produce a spray of high surface-

to-mass ratio droplets, and then to uniformly and quickly evaporate the water.

Evaporation keeps product temperature to a minimum, so little high-temperature deterioration occurs.

The feed can be a solution, a suspension or a paste in the simplest
5 form. The dried product can be powdered, granulated or agglomerated. The dry
product form can be varied depending on the feed, dryer design and process
conditions. Spray drying delivers a powder of specific particle size and moisture
content regardless of the dryer capacity or product heat sensitivity. In a continuous
operation, the spray-dryer delivers a highly controlled powder quality with relatively
10 easy control.

Topping: As used herein, the term "toppings" is taken to include, for
example, salad dressing, whipped toppings, sauce, soy sauce and gravy.

Stability: As used herein, the term "stability" refers to the ability of
the fiber particulate to resist sedimentation and/or stratification which occurs due to
15 separation of the fiber particles from the main liquid suspension. Thus, a fiber
particulate or a product comprising the fiber particulate that is characterized as
"stable" is one in which the fiber particles remain suspended in the main liquid
suspension for the shelf life of the product.

2. Processes for Preparing a Soya Fiber Particulate

20 Various sources of soya fiber may be used to obtain the soya fiber
particulate of the present invention. Suitable soya include but is not limited to whole
soybeans, full fat soy flour, full fat soy flakes, partially defatted soy flour, partially
defatted soy flakes, defatted soy flour, defatted soy flakes, refatted soy flour, refatted
soy flakes, soy protein concentrate and mixtures thereof.

The disclosed invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 90 microns, wherein at least about 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 85 microns, wherein at least about 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 80 microns, wherein at least about 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 75 microns, wherein at least about 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 70 microns, wherein at least about 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 65 microns, wherein at least about 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of

about 0.01 microns to about 60 microns, wherein at least about 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 55 microns, wherein at least 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 50 microns, wherein at least about 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 45 microns, wherein at least about 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 40 microns, wherein at least about 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 35 microns.

In one embodiment of the invention there is provided a soya fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 90% of said soya fiber particulate has a size in a range of about 0.01 microns to about 75 microns. Another

embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 80% of said soya fiber particulate has a size in a range of about 0.01 microns to about 65 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 70% of said soya fiber particulate has a size in a range of about 0.01 microns to about 55 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 60% of said soya fiber particulate has a size in a range of about 0.01 microns to about 45 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 40% of said soya fiber particulate has a size in a range of about 0.01 microns to about 25 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 30% of said soya fiber particulate has a size in a range of about 0.01 microns to about 15 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 20% of said soya fiber particulate has a size in a range of about 0.01 microns to about 10 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 10% of said soya fiber particulate has a size in a range of about 0.01 microns to about 5 microns.

The disclosed invention also provides a soya fiber particulate having a particle size in a range of about 0.04 microns to about 100 microns, wherein at least

75% to about 90% of said soya fiber particulate has a size in a range of about 0.04 to about 35 microns. The disclosed invention also provides processes for preparing a soya fiber particulate. Furthermore, the invention conveniently enables the practitioner to prepare products comprising the soya fiber particulate of a very smooth texture and of a high nutritional quality.

In one embodiment of the invention there is provided a soya fiber particulate having a particle size in a range of about 0.04 microns to about 100 microns, wherein at least 75% to about 90% of said soya fiber particulate has a size in a range of about 0.04 to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.04 microns to about 90 microns, wherein at least 75% to about 90% of said soya fiber particulate has a size in a range of about 0.04 to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.04 microns to about 85 microns, wherein at least 75% to about 90% of said soya fiber particulate has a size in a range of about 0.04 to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.04 microns to about 80 microns, wherein at least 75% to about 90% of said soya fiber particulate has a size in a range of about 0.04 to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.04 microns to about 75 microns, wherein at least 75% to about 90% of said soya fiber particulate has a size in a range of about 0.04 to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.04 microns to about 70 microns, wherein at least 75% to about 90% of said soya fiber particulate has a size in a range of about 0.04 to about 35 microns. Another

embodiment of the invention provides a soya fiber particulate having a particle size in range of about 0.04 microns to about 65 microns, wherein at least 75% to about 90% of said soya fiber particulate has a size in a range of about 0.04 to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a

5 particle size in a range of about 0.04 microns to about 60 microns, wherein at least 75% to about 90% of said soya fiber particulate has a size in a range of about 0.04 to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.04 microns to about 55 microns, wherein at least 75% to about 90% of said soya fiber particulate has a size in

10 a range of about 0.04 to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.04 microns to about 50 microns, wherein at least 75% to about 90% of said soya fiber particulate has a size in a range of about 0.04 to about 35 microns. Another

embodiment of the invention provides a soya fiber particulate having a particle size in

15 a range of about 0.04 microns to about 45 microns, wherein at least 75% to about 90% of said soya fiber particulate has a size in a range of about 0.04 to about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.04 microns to about 40 microns, wherein at least 75% to about 90% of said soya fiber particulate has a size in a range of about 0.04 to

20 about 35 microns. Another embodiment of the invention provides a soya fiber particulate having a particle size in a range of about 0.04 microns to about 35 microns.

In another aspect of the invention, there is provided products comprising the soya fiber particulate which include but are not limited to beverages,

dry mixes, milk products, tofu, miso, toppings, infant food, desserts, snacks, flour products, meat food products, simulated meat food products and pet food products.

In a preferred embodiment of the invention, there is provided a process for preparing a soya fiber particulate. This process comprises grinding dehulled soya
5 to yield a soya fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns, pasteurizing and/or sterilizing said soya fiber particulate, and homogenizing said soya fiber particulate to yield a soya fiber particulate. Preferably, pasteurizing and/or
10 sterilizing the soy fiber particulate is accomplished by heating the soya fiber particulate to a temperature in a range of about 50°C to about 175°C for at least 4 seconds. Also preferably pasteurizing and/or sterilizing the soy fiber particulate is accomplished by heating the soya fiber particulate to a temperature in a range of about 65°C to about 155°C for at least 4 seconds. In one embodiment, the pasteurizing
15 and/or sterilizing of the soya fiber particulate is accomplished prior to homogenizing said soya fiber particulate. In another embodiment, the homogenizing of the soya fiber particulate is accomplished prior to pasteurizing and/or sterilizing said soya fiber particulate.

In a preferred embodiment, the grinding step useful to the process of
20 the present invention yields soya fiber particulate of a median particle size in a range of about 10 microns to about 20 microns. Preferably, at least about 40% of the soya fiber particulate yielded by the grinding step of the present invention has a particle size in a range of about 0.01 microns to about 25 microns. Also preferred, at least about 30% of the soya fiber particulate yielded by the grinding step of the present
25 invention has a particle size in a range of about 0.01 microns to about 15 microns.

Also preferred, at least about 20% of the soya fiber particulate yielded by the grinding step of the present invention has a particle size in a range of about 0.01 microns to about 10 microns. Also preferred, at least about 10% of the soya fiber particulate yielded by the grinding step of the present invention has a particle size in a range of
5 about 0.01 microns to about 5 microns.

In one embodiment, the grinding comprises wet grinding. In another embodiment, the grinding comprises dry grinding. In a specific embodiment, the grinding is performed by a bead mill. In another specific embodiment, the grinding is performed by a ball mill. In a further embodiment, the soya fiber is ground in an
10 apparatus selected from, but in no way limited to, the group consisting of a vibratory mill, an agitator mill, an attrition mill, a roller ball mill, a planetary mill, a hammer mill, a toothed disc mill, a cone mill, a sand mill and a shearing wet mill with a rotor and stator arrangement.

In another preferred embodiment of the invention, there is provided a
15 process for preparing a soya fiber particulate. This process comprises grinding dehulled soybeans to yield a soya fiber particulate having a particle size in a range of about 0.04 microns to about 100 microns, wherein at least 75% to about 90% of said soya fiber particulate has a range of about 0.04 to about 35 microns; heating the soya fiber particulate to a temperature in a range of about 65°C to about 155°C for at least 4
20 seconds; cooling the processed soya fiber particulate to a temperature in a range of about 60°C to about 100°C; and homogenizing the cooled soya fiber particulate to yield a soya fiber particulate.

In a preferred embodiment, the grinding step useful to the process of the present invention yields soya fiber particulate of a median particle size in a range
25 of about 10 microns to about 20 microns. Preferably, at least about 75% to about 90%

of the soya fiber particulate yielded by the grinding step of the present invention has a particle size in a range of about 0.04 microns to about 35 microns. Also preferred, at least about 50% to about 75% of the soya fiber particulate yielded by the grinding step of the present invention has a particle size in a range of about 0.04 microns to about 25 microns. Also preferred, at least about 25% to about 50% of the soya fiber particulate yielded by the grinding step of the present invention has a particle size in a range of about 0.04 microns to about 15 microns. Also preferred, at least about 10% to about 25% of the soya fiber particulate yielded by the grinding step of the present invention has a particle size in a range of about 0.04 microns to about 5 microns.

10 In a specific embodiment, the grinding step is performed by a bead mill. In another specific embodiment, the grinding step is performed by a ball mill. In a further embodiment, the soya fiber ground is an apparatus selected from, but in no way limited to, the group consisting of a vibratory mill, an agitator mill, an attrition mill, a roller ball mill, a planetary mill, a hammer mill, a toothed disc mill, a cone mill, a sand mill and a shearing wet mill with a rotor and stator arrangement.

In another preferred aspect of the invention, there is provided a process for preparing a soya fiber particulate. This process comprises wet grinding dehulled soybeans to yield a soya fiber particulate having a particle size in a range of about 0.04 microns to about 100 microns, wherein at least 75% to about 90% of said soya fiber particulate has a size in a range of about 0.04 to about 35 microns; heating the soya fiber particulate to a temperature in a range of about 65°C to about 155°C for at least 4 seconds; cooling the processed soya fiber particulate to a temperature in a range of about 60°C to about 100°C; and homogenizing the cooled soya fiber particulate to yield a soya fiber particulate.

In a preferred embodiment, the wet grinding step useful to the process of the present invention yields soya fiber particulate of a median particle size in a range of about 10 microns to about 20 microns. Preferably, at least about 75% to about 90% of the soya fiber particulate yielded by the wet grinding step of the present invention has a particle size in a range of about 0.04 microns to about 35 microns. Also preferred, at least about 50% to about 75% of the soya fiber particulate yielded by the wet grinding step of the present invention has a particle size in a range of about 0.04 microns to about 25 microns. Also preferred, at least about 25% to about 50% of the soya fiber particulate yielded by the wet grinding step of the present invention has a particle size in a range of about 0.04 microns to about 15 microns. Also preferred, at least about 10% to about 25% of the soya fiber particulate yielded by the wet grinding step of the present invention has a particle size in a range of about 0.04 microns to about 5 microns.

In a specific embodiment, the wet grinding step is performed by a bead mill. In another specific embodiment, the wet grinding step is performed by a ball mill. In a further embodiment, the soya fiber is ground in an apparatus selected from, but in no way limited to, the group consisting of a vibratory mill, an agitator mill, an attrition mill, a roller ball mill, a planetary mill, a hammer mill, a toothed disc mill, a cone mill, a sand mill and shearing wet mill with a rotor and stator arrangement.

In another preferred aspect of the invention, there is provided a process for preparing a soya fiber particulate. This process comprises dry grinding dehulled soybeans to yield a soya fiber particulate having a particle size in a range of about 100 microns to about 300 microns, wherein at least 75% to about 90% of said soya fiber particulate has a size in a range of about 100 to about 200 microns; heating the soya fiber particulate to a temperature in a range of about 65°C to about 155°C for at least 4

seconds; cooling the processed soya fiber particulate to a temperature in a range of about 60°C to about 100°C; and homogenizing the cooled soya fiber particulate to yield a soya fiber particulate.

In a preferred embodiment, the dry grinding step useful to the process of the present invention yields soya fiber particulate of a median particle size in a range of about 10 microns to about 20 microns. Also preferred, at least about 75% to about 90% of the soya fiber particulate has a particle size in a range of about 0.04 microns to about 35 microns. Also preferred, at least about 50% to about 75% of the soya fiber particulate has a particle size in a range of about 0.04 microns to about 25 microns. Also preferred, at least about 25% to about 50% of the soya fiber particulate has a particle size in a range of about 0.04 microns to about 15 microns. Also preferred, at least about 10% to about 25% of the soya fiber particulate has a particle size in a range of about 0.04 microns to about 5 microns.

In a specific embodiment, the dry grinding step is performed by a bead mill. In another specific embodiment, the dry grinding step is performed by a ball mill. In a further embodiment, the soya fiber is ground in an apparatus selected from, but in no way limited to, the group consisting of a vibratory mill, an agitator mill, an attrition mill, a roller ball mill, a planetary mill, a roller ball mill, a planetary mill, a hammer mill, a toothed disc mill, a cone mill, a sand mill and a shearing wet mill with a rotor and stator arrangement.

In a specific embodiment of the processes for preparing a soya fiber particulate, the soya fiber particulate is heated to a temperature in a range of about 65°C to about 155°C for at least 4 seconds. In another specific embodiment, the soya fiber particulate is heated to a temperature in a range of about 65°C to about 155°C for at least 6 seconds. In another specific embodiment, the soya fiber particulate is

headed to a temperature in a range of about 65°C to about 155°C for at least 10 seconds. In another specific embodiment, the soya fiber particulate is heated to a temperature in a range of about 65°C to about 155°C for at least 15 seconds. In another specific embodiment, the soya fiber particulate is heated to a temperature in a range of about 65°C to about 155°C for at least 20 seconds. In another specific embodiment, the soya fiber particulate is heated to a temperature in a range of about 65°C to about 155°C for at least 30 seconds. In another specific embodiment, the soya fiber particulate is heated to a temperature in a range of about 65°C to about 155°C for at least 45 seconds. In another specific embodiment, the soya fiber particulate is heated to a temperature in a range of about 65°C to about 155°C for at least 60 seconds. In another specific embodiment, the soya fiber particulate is heated to a temperature in a range of about 65°C to about 155°C for at least 5 min. In another specific embodiment, the soya fiber particulate is heated to a temperature in a range of about 65°C to about 155°C for at least 10 min. In another specific embodiment, the soya fiber particulate is heated to a temperature in a range of about 65°C to about 155°C for at least 30 min. In another specific embodiment, the soya fiber particulate is heated to a temperature in a range of about 65°C to about 155°C for at least 1 hr. In another specific embodiment, the soya fiber particulate is heated to a temperature in a range of about 65°C to about 155°C for at least 3 hr.

20 The processes of preparing a soya fiber particulate according to the invention have the further advantage that they are carried out continuously. Alternatively, some of the steps in the processes of preparing a soya fiber particulate according to the invention may be carried out in batches.

In a specific embodiment of the invention, the soya fiber particulate is cooled and packaged. In another specific embodiment, the soya fiber particulate is dehydrated. In another specific embodiment, the soya fiber particulate is spray-dried.

The products comprising the soya fiber particulate of the present invention include fermented soy products, for example, tempheh, miso, soy sauces, natto and fermented tofu and other fermented soymilk products. Additional fermentation steps well known to one of ordinary skill in the art may be carried out in addition to the process of the present invention in order to obtain these fermented products. Alternatively, the products comprising the soya fiber particulate of the present invention include nonfermented soy products, for example, whole soy flour, soymilk and soymilk products, tofu, okara and yuba.

Another advantage of the present invention is the high stability of the soya fiber particulate and products comprising the soya fiber particulate. The small particle size of the soya fiber particulate as well as the grinding in the process of the present invention are believed to enhance the stability of the soya fiber particulate. The grinding involved in the process of the present invention disrupts and opens up the fiber particles such that water or a water based liquid is allowed into the intracellular spaces. This leads to a higher viscosity in the soya fiber particulate, which, along with the small particle sizes of the fibers, provides a soya fiber particulate that is very stable. Preferably, the soya fiber particulate and products comprising the soya fiber particulate remain stable for at least 6 months. Also preferably, the soya fiber particulate and products comprising the soya fiber particulate remain stable for at least 7 months. Also preferably, the soya fiber particulate and products comprising the soya fiber particulate remain stable for at least 8 months. Also preferably, the soya fiber particulate and products comprising the

soya fiber particulate remain stable for at least 9 months. Also preferably, the soya fiber particulate and products comprising the soya fiber particulate remain stable for at least 10 months. Also preferably, the soya fiber particulate and products comprising the soya fiber particulate remain stable for at least 11 months. Also preferably, the soya fiber particulate and products comprising the soya fiber particulate remain stable for at least 1 year.

In a preferred embodiment, the soya fiber particulate and products comprising the soya fiber particulate have a sedimentation of less than about 5% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes. Preferably, the soya fiber particulate and products comprising the soya fiber particulate have a sedimentation of less than about 4% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes. Also preferably, the soya fiber particulate and products comprising the soya fiber particulate have a sedimentation of less than about 3% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes. Also preferably, the soya fiber particulate and products comprising the soya fiber particulate have a sedimentation of less than about 2% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes. Also preferably, the soya fiber particulate and products comprising the soya fiber particulate have a sedimentation of less than about 1% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes. Also preferably, the soya fiber particulate and products comprising the soya fiber particulate have a sedimentation of about 0% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.

Other grains can also be used to obtain a fiber particulate similar to the soya fiber particulate in the present invention. Such grains include but are not limited to flax, sunflower, rice, canola, corn, wheat, rapeseed, and lupin. Other suitable sources of grain fiber include but are not limited to full fat grains, full fat grain flour, full fat grain flakes, partially defatted grain flour, partially defatted grain flakes, defatted grain flour, defatted grain flakes, refatted grain flour, refatted grain flakes, grain protein concentrate and mixtures thereof. The grains can be used in the method of the present invention to produce grain fiber particulates having the same advantages as the soya fiber particulate of the present invention. In one embodiment, the grain fiber particulate has a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 50% to about 100% of said grain fiber particulate has a size in a range of about 0.01 microns to about 35 microns. In another embodiment, the grain fiber particulate has a median particle size in a range of about 10 microns to about 20 microns.

In another aspect of the invention, there is provided products comprising the grain fiber particulate which include but are not limited to beverages, dry mixes, milk products, tofu, miso, toppings, infant food, desserts, snacks, flour products, meat food products, simulated meat food products and pet food products.

Another aspect of the invention also provides a process for preparing the grain fiber particulate, comprising grinding grain seeds to yield a grain fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 50% to about 100% of said grain fiber particulate has a size in a range of about 0.01 to about 35 microns, pasteurizing and/or sterilizing said grain fiber particulate, and homogenizing said grain fiber particulate to yield a grain fiber particulate. Preferably, the grain fiber particulate and products comprising said

grain fiber particulate remain stable for at least 6 months. Also preferably, the grain fiber particulate and products comprising said grain fiber particulate remain stable for at least 1 year. Also preferably, the grain fiber particulate and products comprising said grain fiber particulate have a sedimentation of less than about 5% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes. Also preferably, the grain fiber particulate and products comprising said grain fiber particulate have a sedimentation of about 0% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.

10 The following examples are illustrative only and are not intended to limit the scope of the invention as defined by the appended claims. It will be apparent to those skilled in the art that various modifications and variations can be made in the methods of the present invention without departing from the spirit and scope of the invention. Thus, it is intended that the present invention covers the modifications and
15 variations of this invention provided they come within the scope of the appended claims and their equivalents.

EXAMPLES

20

EXAMPLE 1

Wet milling bean cracks

3500 grams of organic white hylum soybeans were cleaned.

The beans were cracked and dehulled yielding about 3250 grams of

25 bean cracks.

The bean cracks were added to 18,000 grams of 95°C water.

The slurry was run through a Boston Shearpump Model BSP 37-3.

The resulting slurry was then run through two times a Premier Supermill SM 15. at 5.0 gallons per hour with 1.2-1.7 Zirconox beads at a mill speed of 2700 feet per minute.

5 The product was then jet cooked at 115°C for 60 seconds, flash cooled under a pressure of 380 mm of mercury to a temperature of 82°C.

The product was then homogenized at 510 bar on the first stage and 34 bar on the second stage to a total of 544 bar of an APV Rannie 10.38 homogenizer.

10 The product was then spray dried on a tall form dryer with an air inlet temperature of 232°C and an air outlet temperature of 82°C.

Results gave a soymilk product after standard dairy processing that was smooth and without off beany, grassy, or oxidized flavors. The product was smooth in the mouth without being astringent or gritty.

15 The particle sizes of the soya fiber particulates were such that the mean particle size was about 22 microns and 90% of the particles were less than 44 microns.

EXAMPLE 2

Dry milling bean cracks

20 The beans are cracked and dehulled yielding about 3250 grams of bean cracks.

The bean cracks are dry ground with a Bauermeister universal mill and the resulting powder is added to 18,000 grams of 95°C.

The resulting slurry is then run through two times a Premier Supermill SM 1.5 at 5.0 gallons per hour with 1.2-1.7 Zirconox beads at a mill speed of 2700 feet per minute.

The product was then jet cooked at 115°C for 60 seconds, flash cooled under a pressure of 380 mm of mercury to a temperature of 82°C.

The product was then homogenized at 510 bar on the first stage and 34 bar on the second stage to a total of 544 bar of an APV Rannie 10.38 homogenizer.

The product was then sprayed dried on a tall form dryer with an air inlet temperature of 232°C and an air outlet temperature of 82°C.

Results gave a soymilk product after standard dairy processing that was smooth and without off beany, grassy, or oxidized flavors. The product was smooth in the mouth without being astringent or gritty.

The particle sizes of the soya fiber particulates were such that the mean particle size was about 24 microns and 90% of the particles were less than 45 microns.

EXAMPLE 3

Soya fiber particulate particle sizes

Soy milk was made with the soya fiber particulate produced in Example

1. The size of particles in the soy milk was determined, as shown in Table 1.

5

Table 1. Soy milk particle size

% Soya Fiber Particulate	Particle Diameter (microns)
10	3.4
25	8.3
50	17.6
75	31.3
90	46.4
95	57
100	105

EXAMPLE 4

Whole bean soy milk stability and soy fiber particulate stability

- The stability of the soy milk was demonstrated by centrifuging two
- 10 soy milks. The first soy milk is produced as in Example 1, and then reconstituted ultra high temperature pasteurized at 140°C for 4 seconds, homogenized at 170 bar and cooled to 4°C. The second soy milk was a commercial product of similar composition but produced by dry grinding and had a particle size range between 0.5 microns to 350 microns. It was also reconstituted and processed as the first soy milk before
- 15 testing for stability.

The stability test was made using an IEC Centra GP8 centrifuge.

Samples were measured into 50-milliliter centrifuge tubes. The samples were placed in the centrifuge and centrifuged at 500 rpm for 5 minutes.

The results show that the product produced by Example 1 (the first soymilk) does not show a significant amount of sedimentation whereas the commercial product produced by dry grinding (the second soymilk) showed 5% by volume sedimentation.

5 The products were then examined under a light microscope. The product produced by Example 1 showed particles that were smaller and had voids while the dry ground commercial product showed larger particles that were more crystalline in structure.

EXAMPLE 5

10 Defatted soy flakes soymilk stability and soy fiber particulate stability

 The stability of the soymilk was demonstrated by centrifuging a soymilk made with defatted soy flakes using the process described in Example 1. Defatted soy flakes (NUTRISOY 7B Flakes, commercially available from Archer Daniels Midland Company of Decatur, IL) are processed in the same manner as the
15 bean cracks according to Example 1. Specifically, 15,000 grams of defatted soy flakes were mixed with 85,000 grams of 95°C water to produce a slurry. The resulting slurry was then run through two times a Premier Supermill SM 15. at 5.0 gallons per hour with 1.2-1.7 Zirconox beads at a mill speed of 2700 feet per minute. The product was jet cooked at 115°C for 60 seconds, flash cooled under a pressure of
20 380 mm of mercury to a temperature of 82°C. The product was then homogenized at 510 bar on the first stage and 34 bar on the second stage to a total of 544 bar of an APV Rannie 10.38 homogenizer. Next, the product was spray dried on a tall form dryer with an air inlet temperature of 232°C and an air outlet temperature of 82°C. To produce the soymilk product, the dried product was then reconstituted, ultra high

temperature pasteurized at 140°C for 4 seconds, homogenized at 170 bar and cooled to 4°C.

Stability of the soymilk product was measured using an IEC Centra GP8 centrifuge. Samples were deposited into 50-milliliter centrifuge tubes. The
5 samples were placed in the centrifuge and centrifuged at 500 rpm for 5 minutes.

The results show that the product produced by this method with defatted soy flakes does not show a significant amount of sedimentation.

The product was then examined under a light microscope. The product produced with defatted soy flakes by Example 1 showed particles that were small and
10 had voids.

EXAMPLE 6

Flax fiber particulate

Other grains were also run using the procedure of Example 1. For example, 30 pounds whole flax seed was added to 300 pounds of 95°C water. The
15 product was then run through a shear pump which had three grinding heads. A shear pump is a rotor and stator type system which reduces the particle size to approximately 150-200 microns. The product is then run through the bean mill, homogenized, pasteurized and spray dried. Upon reconstitution, the particles' size ranged between 0.04 microns and 95 microns with 80% of the particles being between
20 0.04 microns and 35 microns.

All patents and publications referred to herein are expressly incorporated by reference in their entirety.

WHAT IS CLAIMED IS:

1. A soya fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35
5 microns.
2. The soya fiber particulate of claim 1, wherein at least about 90% of said soya fiber particulate has a particle size in a range of about 0.01 microns to about 75 microns.
3. The soya fiber particulate of claim 2, wherein at least about
10 80% of said soya fiber particulate has a particle size in a range of about 0.01 microns to about 65 microns.
4. The soya fiber particulate of claim 3, wherein at least about 70% of said soya fiber particulate has a particle size in a range of about 0.01 microns to about 55 microns.
- 15 5. The soya fiber particulate of claim 4, wherein at least about 60% of said soya fiber particulate has a particle size in a range of about 0.01 microns to about 45 microns.
6. The soya fiber particulate of claim 1, wherein at least about 40% of said soya fiber particulate has a particle size in a range of about 0.01 microns
20 to about 25 microns.
7. The soya fiber particulate of claim 6, wherein at least about 30% of said soya fiber particulate has a particle size in a range of about 0.01 microns to about 15 microns.

8. The soya fiber particulate of claim 7, wherein at least about 20% of said soya fiber particulate has a particle size in a range of about 0.01 microns to about 10 microns.

9. The soya fiber particulate of claim 8, wherein at least about 10% of said soya fiber particulate has a particle size in a range of about 0.01 microns to about 5 microns.

10. The soya fiber particulate of claim 1, wherein said soya fiber particulate has a median particle size in a range of about 10 microns to about 20 microns.

11. The soya fiber particulate of claim 1, wherein said soya fiber particulate remains stable for at least 6 months.

12. The soya fiber particulate of claim 11, wherein said soya fiber particulate remains stable for at least 7 months.

13. The soya fiber particulate of claim 12, wherein said soya fiber particulate remains stable for at least 8 months.

14. The soya fiber particulate of claim 13, wherein said soya fiber particulate remains stable for at least 9 months.

15. The soya fiber particulate of claim 14, wherein said soya fiber particulate remains stable for at least 10 months.

16. The soya fiber particulate of claim 15, wherein said soya fiber particulate remains stable for at least 11 months.

17. The soya fiber particulate of claim 16, wherein said soya fiber particulate remains stable for at least 1 year.

18. The soya fiber particulate of claim 1, wherein said soya fiber particulate has a sedimentation of less than about 5% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.

19. The soya fiber particulate of claim 18, wherein said soya fiber
5 particulate has a sedimentation of less than about 4% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.

20. The soya fiber particulate of claim 19, wherein said soya fiber particulate has a sedimentation of less than about 3% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.

10 21. The soya fiber particulate of claim 20, wherein said soya fiber particulate has a sedimentation of less than about 2% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.

22. The soya fiber particulate of claim 21, wherein said soya fiber
15 particulate has a sedimentation of less than about 1% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.

23. The soya fiber particulate of claim 22, wherein said soya fiber particulate has a sedimentation of about 0% by volume when centrifuged at a centrifugal force of a least 50 times the force due to the earth's gravity for 5 minutes.

24. The soya fiber particulate of claim 1, wherein said soya is
20 selected from the group consisting of whole soybeans, full fat soy flour, full fat soy flakes, partially defatted soy flour, partially defatted soy flakes, defatted soy flour, defatted soy flakes, refatted soy flour, refatted soy flakes, soy protein concentrate and mixtures thereof.

25. A product comprising the soya fiber particulate of claim 1,
25 wherein said product is selected from the group consisting of:

- 5 (a) beverage;
(b) dry mix;
(c) milk product;
(d) simulated milk product;
(e) tofu;
(f) miso;
(g) topping;
(h) infant food;
(i) dessert;
10 (j) snack;
(k) flour product;
(l) meat food product;
(m) simulated meat food product; and
(n) pet food product.

15 26. A product comprising the soya fiber particulate of claim 1,
wherein said product is soymilk.

27. A product comprising the soya fiber particulate of claim 1,
wherein said product is a simulated milk product combined with a milk product.

28. A product comprising the soya fiber particulate of claim 1,
20 wherein said product is a simulated meat food product combined with a meat food
product.

29. The product of claim 25, wherein said product remains stable
for at least 6 months.

30. The product of claim 29, wherein said product remains stable
25 for at least 7 months.

31. The product of claim 30, wherein said product remains stable
for at least 8 months.
32. The product of claim 31, wherein said product remains stable
for at least 9 months.
- 5 33. The product of claim 32, wherein said product remains stable
for at least 10 months.
34. The product of claim 33, wherein said product remains stable
for at least 11 months.
35. The product of claim 34, wherein said product remains stable
10 for at least 1 year.
36. The product of claim 25, wherein said product has a
sedimentation of less than about 5% by volume when centrifuged at a centrifugal
force of at least 50 times the force due to the earth's gravity for 5 minutes.
37. The product of claim 36, wherein said product has a
15 sedimentation of less than about 4% by volume when centrifuged at a centrifugal
force of at least 50 times the force due to the earth's gravity for 5 minutes.
38. The product of claim 37, wherein said product has a
sedimentation of less than about 3% by volume when centrifuged at a centrifugal
force of at least 50 times the force due to the earth's gravity for 5 minutes.
- 20 39. The product of claim 38, wherein said product has a
sedimentation of less than about 2% by volume when centrifuged at a centrifugal
force of at least 50 times the force due to the earth's gravity for 5 minutes.
40. The product of claim 39, wherein said product has a
sedimentation of less than about 1% by volume when centrifuged at a centrifugal
25 force of at least 50 times the force due to the earth's gravity for 5 minutes.

41. The product of claim 40, wherein said product has a sedimentation of about 0% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.

42. The product of claim 25, wherein said soya is selected from the group consisting of whole soybeans, full fat soy flour, full fat soy flakes, partially defatted soy flour, partially defatted soy flakes, defatted soy flour, defatted soy flakes, refatted soy flour, refatted soy flakes, soy protein concentrate and mixtures thereof.

43. A process for preparing the soya fiber particulate of claim 1, wherein said process comprises:

(a) grinding a dehulled soya to yield a soya fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 50% to about 100% of said soya fiber particulate has a size in a range of about 0.01 microns to about 35 microns;

(b) pasteurizing and/or sterilizing said soya fiber particulate; and

(c) homogenizing said soya fiber particulate to yield a soya fiber particulate.

44. The process of claim 43, wherein said soya is selected from the group consisting of whole soybeans, full fat soy flour, full fat soy flakes, partially defatted soy flour, partially defatted soy flakes, defatted soy flour, defatted soy flakes, refatted soy flour, refatted soy flakes, soy protein concentrate and mixtures thereof.

45. The process of claim 43, wherein said grinding comprises wet grinding.

46. The process of claim 43, wherein said grinding comprises dry grinding.

47. The process of claim 43, wherein said pasteurizing and/or sterilizing comprises heating said soya fiber particulate to a temperature in a range of about 50°C to about 175°C for at least 4 seconds.

48. The process of claim 47, wherein said pasteurizing and/or
5 sterilizing comprises heating said soya fiber particulate to a temperature in a range of about 65°C to about 155°C for at least 4 seconds.

49. The process of claim 43, wherein said grinding yields soya fiber particulate of a median particle size in a range of about 10 microns to about 20 microns.

10 50. The process of claim 43, wherein said grinding yields soya fiber particulate wherein at least about 40% of said soya fiber particulate has a particle size in a range of about 0.01 microns to about 25 microns.

51. The process of claim 50, wherein said grinding yields soya fiber particulate wherein at least about 30% of said soya fiber particulate has a particle
15 size in a range of about 0.01 microns to about 15 microns.

52. The process of claim 51, wherein said grinding yields soya fiber particulate wherein at least about 20% of said soya fiber particulate has a particle size in a range of about 0.01 microns to about 10 microns.

53. The process of claim 52, wherein said grinding yields soya
20 fiber particulate wherein at least about 10% of said soya fiber particulate has a particle size in a range of about 0.01 microns to about 5 microns.

54. The process of claim 43, wherein said grinding is performed by a bead mill or a ball mill.

55. The process of claim 43, wherein said fiber particulate is
25 dehydrated.

56. The process of claim 55, wherein said fiber particulate is spray-dried.

57. The soya fiber particulate made by the process of any of claims 43-56.

5 58. A product comprising the soya fiber particulate of claim 57, wherein said product is selected from the group consisting of:

(a) beverage;

(b) dry mix;

(c) milk product;

10 (d) simulated milk product;

(e) tofu;

(f) miso;

(g) topping;

(h) infant food;

15 (i) dessert;

(j) snack;

(k) flour product;

(l) meat food product;

(m) simulated meat food product; and

20 (n) pet food product.

59. A product comprising the soya fiber particulate of claim 57, wherein said product is soymilk.

60. A product comprising the soya fiber particulate of claim 57, wherein said product is a simulated milk product combined with a milk product.

61. A product comprising the soya fiber particulate of claim 57, wherein said product is a simulated meat food product combined with a meat food product.
62. The product of claim 58, wherein said product remains stable for at least 6 months.
- 5 63. The product of claim 62, wherein said product remains stable for at least 7 months.
64. The product of claim 63, wherein said product remains stable for at least 8 months.
65. The product of claim 64, wherein said product remains stable for at least 9
10 months.
66. The product of claim 65, wherein said product remains stable for at least 10 months.
67. The product of claim 66, wherein said product remains stable for at least 11 months.
- 15 68. The product of claim 67, wherein said product remains stable for at least 1 year.
69. The product of claim 58, wherein said product has a sedimentation of less than about 5% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.
- 20 70. The product of claim 69, wherein said product has a sedimentation of less than about 4% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.
71. The product of claim 70, wherein said product has a sedimentation of less than about 3% by volume when centrifuged at a centrifugal force of at least 50 times the
25 force due to the earth's gravity for 5 minutes.

72. The product of claim 71, wherein said product has a sedimentation of less than 2% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.

73. The product of claim 72, wherein said product has a sedimentation of less than about 1% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.

74. The product of claim 73, wherein said product has a sedimentation of about 0% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.

75. The product of claim 58, wherein said soya is selected from the group consisting of whole soybeans, full fat soy flour, full fat soy flakes, partially defatted soy flour, partially defatted soy flakes, defatted soy flour, defatted soy flakes, refatted soy flour, refatted soy flakes, soy protein concentrated and mixtures thereof.

76. A grain fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 50% to about 100% of said grain fiber particulate has a size in a range of about 0.01 microns to about 35 microns.

77. The grain fiber particulate of claim 76, wherein said grain is selected from the group consisting of flax, sunflower, rice, canola, corn, wheat, rapeseed, and lupin.

78. The grain fiber particulate of claim 76, wherein said grain is selected from the group consisting of full fat grains, full fat grain flour, full fat grain flakes, partially defatted grain flour, partially defatted grain flakes, defatted grain flour, defatted grain flakes, refatted grain flour, refatted grain flakes, grain protein concentrate and mixtures thereof.

79. The grain fiber particulate of claim 76, wherein said grain fiber particulate has a median particle size in a range of about 10 microns to about 20 microns.

80. The grain fiber particulate of claim 76, wherein said grain fiber particulate remains stable for at least 6 months.
81. The grain fiber particulate of claim 80, wherein said grain fiber particulate remains stable for at least 1 year.
- 5 82. The grain fiber particulate of claim 76, wherein said grain fiber particulate has a sedimentation of less than about 5% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.
83. The grain fiber particulate of claim 82, wherein said grain fiber particulate has a sedimentation of about 0% by volume when centrifuged at a centrifugal force of at
- 10 least 50 times the force due to the earth's gravity for 5 minutes.
84. A product comprising the grain fiber particulate of claim 76, wherein said product is selected from the group consisting of:
- (a) beverage;
 - (b) dry mix;
 - 15 (c) milk product;
 - (d) simulated milk product;
 - (e) tofu;
 - (f) miso;
 - (g) topping;
 - 20 (h) infant food;
 - (i) dessert;
 - (j) snack;
 - (k) flour product;
 - (l) meat food product;
 - 25 (m) simulated meat food product; and

(n) pet food product.

85. The product of claim 84, wherein said grain is selected from the group consisting of flax, sunflower, rice, canola, corn, wheat, rapeseed, and lupin.

86. The product of claim 84, wherein said grain is selected from the group
5 consisting of full fat grains, full fat grain flour, full fat grain flakes, partially defatted grain flour, partially defatted grain flakes, defatted grain flour, defatted grain flakes, refatted grain flour, refatted grain flakes, grain protein concentrate and mixtures thereof.

87. The product of claim 84, wherein said product remains stable for at least 6
10 months.

88. The product of claim 87, wherein said product remains stable for at least 1 year.


89. The product of claim 84, wherein said product has a sedimentation of less than about 5% by volume when centrifuged at a centrifugal force of at least 50 times the
15 force due to the earth's gravity for 5 minutes.

90. The product of claim 89, wherein said product has a sedimentation of about 0% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.

91. A process for preparing the grain fiber particulate of claim 76, wherein said
20 process comprises:

(a) grinding grain seeds to yield a grain fiber particulate having a particle size in a range of about 0.01 microns to about 100 microns, wherein at least about 50% to about 100% of said grain fiber particulate has a size in a range of about 0.01 to about 35 microns;

25 (b) pasteurizing and/or sterilizing said grain fiber particulate; and



(c) homogenizing said grain fiber particulate to yield a grain fiber particulate.


92. The process of claim 91, wherein said grain is selected from the group consisting of flax, sunflower, rice, canola, corn, wheat, rapeseed, and lupin.

5 93. The process of claim 91, wherein said grain is selected from the group consisting of full fat grains, full fat grain flour, full fat grain flakes, partially defatted grain flour, partially defatted grain flakes, defatted grain flour, defatted grain flakes, refatted grain flour, refatted grain flakes, grain protein concentrate and mixtures thereof.

10 94. The grain fiber particulate made by the process of any of claims 91-93.

95. A product comprising the grain fiber particulate of claim 94, wherein said product is selected from the group consisting of:

- (a) beverage;
- (b) dry mix;
- 15 (c) milk product;
- (f) simulated milk product;
- (d) tofu;
- (e) miso;
- (f) topping;
- 20 (g) infant food;
- (h) dessert;
- (i) snack;
- (j) flour product;
- (k) meat food product;
- 25 (l) simulated meat food product; and



(m) pet food product.

96. The product of claim 95, wherein said grain is selected from the group consisting of flax, sunflower, rice, canola, corn, wheat, rapeseed, and lupin.

97. The product of claim 95, wherein said grain is selected from the group
5 consisting of full fat grains, full fat grain flour, full fat grain flakes, partially defatted grain flour, partially defatted grain flakes, defatted grain flour, defatted grain flakes, refatted grain flour, refatted grain flakes, grain protein concentrate and mixtures thereof.

98. The product of claim 95, wherein said product remains stable for at least 6
10 months.

99. The product of claim 98, wherein said product remains stable for at least 1 year.

100. The product of claim 95, wherein said product has a sedimentation of less than about 5% by volume when centrifuged at a centrifugal force of at least 50 times the
15 force due to the earth's gravity for 5 minutes.

101. The product of claim 100, wherein said product has a sedimentation of about 0% by volume when centrifuged at a centrifugal force of at least 50 times the force due to the earth's gravity for 5 minutes.